

UNCLASSIFIED

AD NUMBER

AD014838

CLASSIFICATION CHANGES

TO: **unclassified**

FROM: **secret**

LIMITATION CHANGES

TO:
Approved for public release, distribution unlimited

FROM:
Distribution authorized to DoD and DoD contractors only; Foreign Gov't. Info.; 15 Jul 1953. Other requests shall be referred to the British Embassy, 3100 Massachusetts Ave., NW, Washington, DC 20008.

AUTHORITY

Public Record Number WO 189/699; Public Record Number WO 189/699

THIS PAGE IS UNCLASSIFIED

UNCLASSIFIED


AD _____

DEFENSE DOCUMENTATION CENTER

FOR

SCIENTIFIC AND TECHNICAL INFORMATION

CAMERON STATION ALEXANDRIA, VIRGINIA



UNCLASSIFIED

Reproduced by

Armed Services Technical Information Agency
DOCUMENT SERVICE CENTER

KNOTT BUILDING, DAYTON, 2, OHIO

AD -

1 4 8 3 8

BEST

AVAILABLE

COPY

14838
SECRET

P.T.P. 365

P.T.P. 365

Copy No. 66

MINISTRY OF SUPPLY

DIRECTORATE OF CHEMICAL DEFENCE RESEARCH AND DEVELOPMENT

CHEMICAL DEFENCE EXPERIMENTAL ESTABLISHMENT

THE EFFECT OF GB ON THE RAT'S BLOOD PRESSURE

By

P. DIRNHUBER AND H. CULLUMBINE

PORTON TECHNICAL PAPER No. 365

C.D.E.E.
Porton.
Wilts.

SECRET

5341 - 11328

SECRET

PORTON TECHNICAL PAPER NO. 365

Copy No. 66

Date 15 JUL 1955

The Effect of GB on the rat's blood pressure

by

P. Dirnhuber and H. Cullumbine

SUMMARY

GB, DFP, eserine, TBT and E.600 all produce hypertension when administered to rats in near-lethal doses. The mechanism of production of this hypertension has been studied. It is, apparently, due to a central mechanism acting via the sympathetic nervous system on the blood vessels of the skin.

(Sgd.) H. Cullumbine,
Head, Physiology Section.

(Sgd.) E.A. Porron,
Superintendent, Research Division.

PD/HC/ST

SECRET

SECRET

PORTON TECHNICAL PAPER NO. 365

Copy No. 66

Date 11 Nov 1954

The Effect of GB on the rat's blood pressure

by

P. Dirnhuber and H. Cullumbine

Introduction

In recent species studied (e.g. rabbit, cat, monkey, dog) the effect of a systemic intoxication with GB is the production of a profound fall in blood pressure. Wilson has analysed the mechanisms responsible for this hypotensive action.

We have recently noticed that GB, when given intravenously in near lethal doses (40 - 60 $\mu\text{g}/\text{kg.}$) to intact rats, produces a sharp rise in blood pressure, then one or two oscillations about the point of increased pressure, followed by a very gradual fall in pressure over the next several minutes (10 - 180 minutes in different animals) back to the pre-injection level (Figure 1). The course of this sustained rise in blood pressure produced by GB has been investigated.

Methods

White rats of homogenous strain and weighing 350 - 500 g. were used. They were anaesthetised with urethane (1.25g./kg. subcutaneously) and polythene canulae inserted into the carotid artery, to record blood pressure, and into the femoral vein.

With the doses of GB used respiratory embarrassment or failure may occur and may interfere with the blood pressure recording or response. Therefore, in many cases, even respiratory exchange was maintained throughout by means of a miniature starting "Ideal" pump. The same pattern of cardiovascular response has, however, been seen in rats which were not sustained by artificial ventilation.

Spinal preparations were made by transecting the spinal cord between C1 and C2 and pithing the brain.

Results:- The hypertensive effect of GB is best seen following a fairly large dose (40 - 60 $\mu\text{g}/\text{kg.}$) but is still evident with smaller doses. The effects of rapidly repeated small doses can be summated up to a certain point and then the general blood pressure level falls although each

SECRET

SECRET

successive GB dose produces a transient, small rise (Figure 2). With higher doses of GB (e.g. 90 $\mu\text{g/kg}$) the animal dies before the sustained rise in blood pressure has become established (Figure 3).

Bilateral vagotomy does not affect the picture (Figure 4), but if GB is given to a spinal rat only a relatively slow, small and short lasting increase in blood pressure is caused (Figure 5).

The failure of GB to produce a sustained elevation of blood pressure in this case is not due to the low blood pressure presented by a spinal rat. Thus, if the blood pressure is lowered by bleeding (Figure 6) or by O_6 (Figure 10). GB still produces a typical hypertensive blood pressure response.

This suggests that GB is stimulating a central mechanism and this stimulation is probably a cholinergic phenomenon since in the previously atropinized cat (intact or spinal) GB produces again only a slow and minor rise in blood pressure (Figure 7). Similarly, if successive doses of atropine are administered after the GB, a step-like depression of the elevated blood pressure is produced (Figure 8).

If hexamethonium bromide (O_6) is given to a rat during the phase of sustained blood pressure rise following GB, the blood pressure is immediately and temporarily reduced (Figure 9). Pre-treatment with O_6 before administering the GB does not prevent the usual blood pressure response to the latter (Figure 10).

The central stimulant action of GB is effected, therefore, via the sympathetic nervous system. The liberation of adrenaline from the adrenal glands would not seem to be important since GB still causes an increased blood pressure in adrenalectomized rats (Figure 11) and this can be inhibited by O_6 .

A sympathetic peripheral vascular mechanism is probably involved since peripheral blockade of sympathetic impulses with ergotamine or Priscol does alter the response to GB. If either of these substances is administered before GB, the latter produces only a preliminary sharp rise, but no sustained elevation of blood pressure (Figure 12). If GB is given first and then Priscol during the period of raised blood pressure, the latter is at once temporarily reduced (Figure 13).

The sustained rise in blood pressure induced by GB is probably due mainly to constriction of the skin arterioles; in the skinned rat, GB causes only an immediate and temporary rise in pressure, but no continued elevation is seen (Figure 14). In conformity with this is the observation that GB produces a maintained pressure rise when administered to an eviscerated rat (Figure 15). This rise is not as long lasting as that occurring in the intact rat, so that some involvement of the arterioles of the splanchnic area may be present.

That a direct action on a central mechanism is involved is also suggested by the observation that a small dose (5 μg) of GB, injected into the fourth ventricle, causes a similar sustained rise in blood pressure (Figure 16). An injection of acetyl choline into the same site

SECRET

produces a fall in blood pressure (Figure 17). The dissimilarity in the actions of acetyl choline and GB may be due to a failure of the former to penetrate into the brain substance, although some leakage into the systemic circulation occurs. The direct central action of the GB can still be prevented by systemic atropinization (Figure 18).

The hypertensive action of GB in the rat is not peculiar to that compound, but is shared by other anti-cholinesterases, e.g. DFP (Figure 19), eserine, TEPP, E.600, and probably many others. Atropine and OG will affect the response to these agents in the manner described for GB.

Another indication of the enhanced sympathetic tone in these rats is that the heart rate (measured via the E.C.G.) is increased during the period of hypertension e.g. in a typical instance:

Heart rate before GB (40 µg/kg.) - 270/minute

Heart rate during phase of rising blood pressure - 390/minute

Heart rate at peak of blood pressure rise - 450/minute

In other species where a fall of blood pressure is produced by GB, this is accompanied by a slowing of the heart rate (Wilson (1)).

Discussion:- From the above results we can conclude that GB, when given intravenously to the rat, produces a rise in blood pressure which is due to a central action, possibly via the vasomotor centre. The effect is not seen in the spinal rat, nor following atropine. The latter observation suggests that the action is probably a cholinergic phenomenon involving the centre.

The central stimulation presumably acts via the sympathetic nervous system and this sympathetic "drive" can be blocked at the ganglia by hexamethonium bromide and also more peripherally by ergotamine or Priscol. It would appear that the vessels of the rat's skin are those chiefly concerned in this sympathetic action.

Other anticholinesterases can produce a similar hypertension in the rat, so that the phenomenon presumably has the inhibition of cholinesterase as its basis. The rise in blood pressure is, further, not peculiar to the intravenous route of administration since it has also been seen after the intraventricular or the intracerebral injection of GB.

In other species GB, when administered intravenously in near-lethal doses, causes a profound lowering of the blood pressure. The latter is accompanied by a marked slowing of the heart and vasodilatation of the small vessels in the limb muscles (Wilson, (1); Holmes (2)). If, however, GB in small doses is injected into the vertebral artery or the cisterna magna of a dog, a rise in blood pressure is the invariable response (Wilson, (3)). Therefore a hypertensive response via a central mechanism can be seen in another species than the rat. In the dog, following intravenous administration, it must be presumed that the peripheral effects of GB on the cardiovascular system predominate over the central stimulating action, although a transient rise of B.P. is often observed before the profound fall takes place. This applies equally to the sheep (Wilson (4)).

SECRET

These varying responses in the different species suggest that there may be corresponding differences in the nature and the sensitivity of the receptor substances and the cholinesterases of the tissues of these species. This is being further investigated.

Summary

GB, DFP, eserine, TEPP and E.600 all produce hypertension when administered to rats in near-lethal doses. The mechanism of production of this hypertension has been studied. It is, apparently, due to a central mechanism acting via the sympathetic nervous system on the blood vessels of the skin.

(Sgd.) H. Cullumbino,
Head, Physiology Section.

(Sgd.) E.A. Perron,
Superintendent, Research Division.

References

- (1) Wilson, K.M. P.T.P. 152.
- (2) Holmes, R. P.T.P. 356.
- (3) Wilson, K.M. Personal communication.
- (4) Wilson, K.M. and Lärnhuber, P. P.T.P. 346.

- 4 -

SECRET




FIG.1. HYPERTENSIVE EFFECT OF GB.

Rat 370g. Urethane.

Spontaneous respiration
at arrow 40 μ GB/kg intravenously.




FIG.3. HIGHER DOSE OF GB.

Rat 420g. Urethane.
90 μ GB/kg intravenously at
arrow 1. Interrupted artificial
ventilation after arrow 2.




FIG.2. REPEATED SMALL DOSES OF GB

Rat 400g. Urethane. Artificial ventilation.

At each arrow 5 μ GB/kg intravenously.
(Total of 11 doses)

T936/1

Time signal: 1 minute intervals

1-2




FIG. 4. RESPONSE NOT AFFECTED BY VAGOTOMY.

Rat 400g. Urethane. Both vagi cut, carotid artery tied. Artificial ventilation.
At arrow 40 μ g GB/kg intravenously.




FIG. 5. SMALL RISE IN SPINAL RAT.

Rat 470g Urethane. Spine trans sectioned, brain pithed.
At arrow 40 μ g GB/kg intravenously.




FIG. 6. TYPICAL PRESSOR RESPONSE AFTER EXSANGUINATION.

Rat 470g Urethane. Artificial ventilation
6ml. blood withdrawn between strip 1 & 2.
At arrow 40 μ g GB/kg intravenously.




FIG. 7. SMALL RISE IN ATROPINISED ANIMAL
Rat 310g Urethane
Artificial ventilation
At arrow 1 10mg ATR/kg
At arrow 2 40 μ g GB/kg intravenously.




FIG. 8. ATROPINE LOWERS ELEVATED BP
Rat 420g. Urethane. Artificial ventilation.
At arrow 1 $40\mu\text{g CB/kg}$ i.v.
At arrows 2 to 5 each 0.5mg ATR/kg i.v.




FIG. 9. C6 AFTER CB
REDUCES BP
Rat 390g. Urethane.
Artificial ventilation.
At arrow 1 $40\mu\text{g CB/kg}$.
At arrow 2 10mg C6/kg .
intravenously.




FIG. 10. PRE-TREATMENT WITH C6.
TYPICAL PRESSOR RESPONSE BY CB
Rat 390g. Urethane. Spontaneous respiration
At arrow 1 20mg C6/kg } intravenously.
2 $60\mu\text{g C6/kg}$ }




FIG. 11.
PRESSOR RESPONSE
UNAFFECTED BY
ADRENALECTOMY.
Rat 480g. Urethane.
Artificial ventilation.
Both adrenals removed.
At arrow $40\mu\text{g CB/kg}$
intravenously. T. 9/3/5.

Time signal 1 minute intervals




FIG.12. PRE-TREATMENT WITH PRISCOL
SMALL PRESSOR EFFECT

Rat 390g Urethane
Artificial ventilation
At arrow 1 10mgPRI/kg
At arrow 2 40µgGB/kg




FIG.13. PRISCOL LOWERS ELEVATED BP

Rat 470g Urethane Artificial ventilation
At arrow 1 40µgGB/kg
At arrows 2 to 5 each 1 mg PRI/kg
all intravenously




FIG.14 NO CONTINUED ELEVATION IN SKINNED ANIMAL

Rat 480g Urethane Artificial ventilation
All skin removed (head, feet & scrotum not skinned)
Body placed in saline bath of 37°
At arrow 40µgGB/kg intravenously.




FIG.15. MAINTAINED ELEVATION IN EVISCERATED ANIMAL.

Rat 390g. Urethane.
Artificial ventilation
Small and large intestines removed.
At arrow 40µgGB/kg i.v.

T.936/4

Time signal: 1 minute intervals




FIG. 16. SMALL DOSE OF CB INTO 4th VENTRICLE
Rat 420g. Urethane. Artificial ventilation.
At arrow 5 µg CB/rat through atlanto-occipital
membrane




FIG. 17. OPPOSITE EFFECT OF
ACh AND GB ON INJECTION
INTO 4th VENTRICLE
Rat 370g. Urethane.
Artificial ventilation.
At arrow 1 2 µg ACh/rat
At arrow 2 5 µg GB/rat
both through atlanto-occipital
membrane.




FIG. 18. PRE TREATMENT WITH
ATROPINE INHIBITS PRESSOR
RESPONSE TO INTRAVENTRICULAR GB
Rat 350g. Urethane. Artificial ventilation.
At arrow 1 2 µg ACh/kg
2 2 mg ATR/kg } intravenously.
3 2 mg ACh/kg
4 5 µg GB/rat into ventricle.




FIG. 19. HYPERTENSIVE EFFECT OF DFP.
Rat 470g. Urethane. Artificial ventilation.
At arrow 1 mg DFP/kg
intravenously.

T. 924/5.

Time signal 1 minute intervals

P.T.P. 365

CIRCULATION
INTERNAL

Copy No.

1 P.D.S.R.(D)
2 D.C.D.R.D.
26 - 115 C.S., C.D.E.E.
3 - 4 C.D.R. Branches
5 - 6 C.S., M.O.S. Nancekuke
7 D.P.B.R.
8 R.D.Arm 2.
9 - 10 T.P.A.3/T.I.B.
11 - 25 S.A.C. for Biology Committee,
C.D.A.B.
26 File

Chemical Defence Advisory Board

27 Professor Sir Rudolph Peters
28 Professor G.R. Cameron
29 J. Davidson Pratt, Esq.
30 Professor H.J. Emelius
31 Sir Paul Fildes
32 Professor J.H. Gadnam
33 Professor Sir Robert Robinson
34 Professor R.H.S. Thompson

EXTERNAL

British Joint Services Mission

41 - 52 D.C. Evans, Esq.

War Office

53 S.W.V.1.(b)

Copy No.

Biology Committee

35 Dr. J.M. Barnes
36 Professor I de Borch Daly
37 Dr. G.S. Dawer
38 Dr. Malcolm Dixon
39 Dr. Bentley Purchase
40 Professor A. Wilson

OVERSEAS

(through T.P.A. 3/T.T.P.)

Australia

54 - 56 Defence Research Laboratories
57 Senior Representative, Dept. of
Supply
58 Army Branch Representative
59 R.A.A.F. (Tech. Section)

Canada

60 - 61 Chairman, Defence Research Board
62 - 63 Defence Research Laboratories,
Ottawa
64 Suffield Experimental Station

U.S.A.

65 - 82 Reading Committee

Reproduced by

Armed Services Technical Information Agency
DOCUMENT SERVICE CENTER

KNOTT BUILDING, DAYTON, 2, OHIO

AD -

1 4 8 3 8